

AD-A056 776

PURDUE UNIV LAFAYETTE IND DEPT OF STATISTICS F/G 12/1  
AN ESSENTIALLY COMPLETE CLASS OF MULTIPLE DECISION PROCEDURES.(U)  
JUN 78 S S GUPTA, D HUANG N00014-75-C-0455  
MMS-78-12 NL

UNCLASSIFIED

| OF |  
AD  
A056776



END  
DATE  
FILMED  
9-78  
DDC


AD No.            AD A056776  
DDC FILE COPY

LEVEL *III*

*(D)*  
*B.S.*



# PURDUE UNIVERSITY



## DEPARTMENT OF STATISTICS

### DIVISION OF MATHEMATICAL SCIENCES

This document has been approved  
for public release and sale; its  
distribution is unlimited.

78 07 24 028

(1)

(6) An Essentially Complete Class of Multiple  
Decision Procedures<sup>†</sup>.

(10) by  
Shanti S./Gupta Purdue University  
and  
Deng-Yuan/Huang Academia Sinica, Taipei, Taiwan

(12) 12 p.

(9) Mimeograph series,

Department of Statistics  
Division of Mathematical Sciences  
Mimeograph No. 78-12

DDC  
RECEIVED  
JUL 28 1978  
F

(11) June 1978

(14) MMS-78-12

(15) \*This research was supported by the Office of Naval Research Contract  
N00014-75-C-0455 at Purdue University. Reproduction in whole or in part  
is permitted for any purpose of the United States Government.

This document has been approved  
for public release and sale; its  
distribution is unlimited.

78 07 25 028  
292 730

set



# Abstract

Let  $\pi_1, \dots, \pi_k$  represent  $k$  ( $\geq 2$ ) independent populations. The quality of the  $i$ th population  $\pi_i$  is characterized by a real-valued parameter  $\theta_i$ , usually unknown. We define the best population in terms of a measure of separation between  $\theta_i$ 's. A selection of a subset containing the best population is called a correct selection (CS). We restrict attention to rules for which the size of the selected subset is controlled at a given point and the infimum of the probability of correct selection over the parameter space is maximized. The main theorem deals with construction of an essentially complete class of selection rules of the above type. Some classical subset selection rules are shown to belong to this class.

## Key Words

Subset selection procedure, monotone likelihood ratio, monotone selection rule, normal means problem, unequal sample sizes.

ACCESSION for		
NTIS	White Section	<input checked="" type="checkbox"/>
DDC	Buff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL.	SPECIAL
A		

An Essentially Complete Class of Multiple  
Decision Procedures\*

by

Shanti S. Gupta, Purdue University

and

Deng-Yuan Huang, Academia Sinica, Taipei, Taiwan

During the past decade, selection and ranking theory has developed rapidly. Many reasonable rules have been proposed. Some good properties have been studied. However, very little work has been done to consider the optimality of a selection procedure, especially in the subset selection approach. In this paper, we discuss an essentially complete class of subset selection rules (in some sense). Some classical selection rules are shown to be optimal in this sense.

Let  $\pi_1, \dots, \pi_k$  represent  $k$  ( $\geq 2$ ) independent populations and let  $x_{i1}, \dots, x_{in_i}$  be  $n_i$  independent random observations from  $\pi_i$ . The quality of the  $i$ th population  $\pi_i$  is characterized by a real-valued parameter  $\theta_i$ ; usually unknown. Let  $\Omega = \{\underline{\theta} | \underline{\theta} = (\theta_1, \dots, \theta_k)\}$  denote the whole parameter space. Let  $\tau_{ij} = \tau_{ij}(\underline{\theta})$  be a measure of separation between  $\tau_i$  and  $\tau_j$ . We assume that there exists a monotone non-increasing function  $h$  such that  $\tau_{ji} = h(\tau_{ij})$ . Let  $\Omega_i = \{\underline{\theta} | \tau_{ij} \geq \tau_{ii}, j \neq i\}$ ,  $1 \leq i \leq k$ . In this sequel, we assume  $\tau_{ii}$  as known,  $1 \leq i \leq k$ . Let  $\tau_i = \min_{j \neq i} \tau_{ij}$ ,  $i = 1, 2, \dots, k$ . Assume that there exists an  $i$  such that  $\tau_i \geq \tau_{ii}$ . Thus we know that  $\Omega = \bigcup_{i=1}^k \Omega_i$ . We define  $\tau^* = \max_{\ell} \tau_{\ell}$ . The population associated with  $\tau^*$  will be called the best population. We know that if  $\underline{\theta} \in \Omega_i$ , then  $\pi_i$  is

---

\*This research was supported by the Office of Naval Research Contract N00014-75-C-0455 at Purdue University. Reproduction in whole or in part is permitted for any purpose of the United States Government.

the best population. A selection of a subset containing the best population is called a correct selection (CS).

We will restrict attention to those selection procedures which depend upon the observations only through a sufficient and maximal invariant statistic  $Z_{ij}$  which is based on the  $n_i$  and  $n_j$  observations from  $\pi_i$  and  $\pi_j$  ( $i, j=1, 2, \dots, k$ ), respectively. It is well known that the distribution of  $Z_{ij}$  depends only on  $\tau_{ij}$ . For any  $i$ , let the joint density of  $Z_{ij}$ ,  $j \neq i$ , be  $p_{\theta}(\underline{z}_i)$ . Let  $p_{\theta}(\underline{z}_i)$  be denoted by  $p_i(\underline{z}_i)$  when  $\tau_{i1} = \dots = \tau_{ik} = \tau_{ii}$ , where  $\underline{z}_i = (z_{i1}, \dots, z_{i,i-1}, z_{i,i+1}, \dots, z_{ik})$ ,  $1 \leq i \leq k$ . Let  $F_{\theta}$  be the continuous cumulative distribution function of  $p_{\theta}(\underline{z})$  for any  $\theta$  and let  $p_i(z_{ij}|z_{i\ell}, \ell \neq i, j)$  be the conditional pdf of  $Z_{ij}$ , given  $Z_{i\ell} = z_{i\ell}$ ,  $\ell \neq i, j$  and let  $F_i(z_{ij}|z_{i\ell}, \ell \neq i, j)$  be the cdf of the conditional density  $p_i(z_{ij}|z_{i\ell}, \ell \neq i, j)$ . Let  $F_i^{\circ}(y)$  be any point of the set  $\{z: F_i(z|z_{i\ell}, \ell \neq i) = y\}$ .

Let  $\delta = (\delta_1, \dots, \delta_k)$  be a selection procedure where  $\delta_i(\underline{z})$  is the probability of selecting  $\pi_i$ ,  $1 \leq i \leq k$ , having observed  $\underline{z}$ . Let  $S(\theta, \delta) = P(\text{CS}|\delta)$  and  $R(\theta, \delta) = \sum_{i=1}^k R^i(\theta, \delta_i)$ , where  $R^i(\theta, \delta_i) = \int \delta_i(\underline{z}_i) p_{\theta}(\underline{z}_i) d\nu(\underline{z}_i)$ . Let  $R_j^i(\delta_i) = \int \delta_i(\underline{z}_i) p_i(z_{ij}|z_{i\ell}, \ell \neq i, j) dz_{ij}$ ,  $1 \leq i \neq j \leq k$ . A decision rule  $\delta_1 = (\delta_{11}, \dots, \delta_{1k})$  is said to be "as good as"  $\delta_2 = (\delta_{21}, \dots, \delta_{2k})$  if  $\inf_{\theta \in \Omega} S(\theta, \delta_1) \geq \inf_{\theta \in \Omega} S(\theta, \delta_2)$  provided that  $\int \delta_{ij} p_j = \gamma_j$ ,  $1 \leq j \leq k$ ,  $i=1, 2$ , where  $\gamma_j$ , ( $0 < \gamma_j < 1$ ), are specified numbers. Let  $C$  be the class of  $\delta$  such that  $\int \delta_i p_i = \gamma_i$ ,  $1 \leq i \leq k$ .

A point  $x_0$  is called a change point for a function  $g$  if in some neighborhood of  $x_0$ ,

$$g(x)g(x^*) \leq 0,$$

whenever  $x \leq x_0 \leq x^*$ , and for some  $x_1 \leq x_0 \leq x_1^*$ ,  $g(x_1) \neq 0$  and  $g(x_1^*) \neq 0$  with  $x_1 \neq x_1^*$ .



Karlin and Rubin [3] have proved the following result.

Lemma ([3]). If  $\varphi$  changes sign at most once in one-dimensional Euclidean space  $R^1$ , then

$$\psi(w) = \int p(x|w) \varphi(x) d\mu(x)$$

changes sign at most once, where  $\mu$  is a  $\sigma$ -finite measure on  $R^1$  and  $p(x|w)$  is the density of  $X$  with monotone likelihood ratio (MLR) in  $w$ .

Remark: It is useful to note that  $\psi$  changes sign in the same direction as  $\varphi$  if it changes sign at all.

Now we define a "monotone" selection rule as follows.

Definition: A selection rule  $\delta$  is called monotone if for any  $i$ ,  $\delta_i(\underline{z})$  is monotone as follows:

$$\delta_i(\underline{z}) = \begin{cases} 1 & \text{if } \underline{z} \geq \underline{z}_0, \\ 0 & \text{otherwise,} \end{cases}$$

where  $\underline{z}_0$  is a fixed known vector and " $\leq$ " is a partial order defined as follows: if  $\underline{x}_1 = (x_{11}, \dots, x_{1m})$  and  $\underline{x}_2 = (x_{21}, \dots, x_{2m})$ , then  $\underline{x}_1 \leq \underline{x}_2 \Leftrightarrow x_{1i} \leq x_{2i}$  for  $i = 1, \dots, m$ .

Theorem: Let  $F_{\underline{\theta}}$  be the continuous cumulative distribution function corresponding to  $p_{\underline{\theta}}(\underline{z})$  which has monotone likelihood ratio. Then all monotone selection procedures form an essentially complete class in  $C$ .

Proof: Let  $\delta$  be any nonmonotone rule in  $C$ . Suppose that there is an  $i$ , such that  $\delta_i$  is not monotone in  $z_{ij}$  for fixed  $z_{i\ell}$ ,  $\ell \neq i, j$ . For each fixed  $z_{i\ell}$  ( $\ell \neq i, j$ ), we define

$$\delta_i^{\circ}(\underline{z}_i) = \begin{cases} 1 & \text{if } z_{ij} \geq f_i^{\circ}(1-R_j^i(\delta_i)), \\ 0 & < \end{cases},$$

then

$$\begin{aligned} (1) \quad \int \delta_i^{\circ} p_i dz_{ij} &= \int_{F_i^{\circ}(1-R_j^i(\delta_i))} p_i dz_{ij} \\ &= \int_{F_i^{\circ}(1-R_j^i(\delta_i))} p_i(z_{ij} | z_{i\ell}, \ell \neq i, j) p(z_{i\ell}, \ell \neq i, j) dz_{ij} = \int \delta_i p_i(\underline{z}_i) dz_{ij}. \end{aligned}$$

Since  $\delta_i^{\circ}$  is monotone in  $z_{ij}$ , thus  $\delta_i - \delta_i^{\circ}$  as a function of  $z_{ij}$  has at most one sign changes from plus to minus. Using this fact, the MLR property of  $p_{\underline{\theta}}(\underline{z}_i)$  and Lemma, we have

$$(2) \quad \int [\delta_i - \delta_i^{\circ}] p_{\underline{\theta}}(\underline{z}_i) dz_{ij} \leq 0, \quad \tau_{ij} \geq \tau_{ii}.$$

Thus from (1) and (2),  $\delta^{\circ}$  has the same conditional size as  $\delta$  and has higher conditional power than  $\delta$  as follows

$$(3) \quad \int \delta_i p_{\underline{\theta}} \leq \int \delta_i^{\circ} p_{\underline{\theta}} \quad \text{for } \tau_{ij} \geq \tau_{ii}, j=1, \dots, k, j \neq i.$$

Since

$$\inf_{\underline{\theta} \in \Omega} S(\underline{\theta}, \delta) = \min_{1 \leq i \leq k} \inf_{\underline{\theta} \in \Omega_i} \int \delta_i p_{\underline{\theta}}(\underline{z}_i) dv(\underline{z}_i),$$

hence by (3),

$$\inf_{\underline{\theta} \in \Omega} S(\underline{\theta}, \delta) \leq \inf_{\underline{\theta} \in \Omega} S(\underline{\theta}, \delta^{\circ}).$$

The proof is complete.



Example: Let  $X_{i1}, \dots, X_{in_i}$  be independent normally distributed with mean  $\theta_i$  and variance  $\sigma^2 = 1$ ,  $i=1, 2, \dots, k$ . Then the joint likelihood function of  $\bar{X}_i$ ,  $1 \leq i \leq k$ , is

$$g_{\underline{\theta}}(\underline{x}) = \prod_{j=1}^k g_{\theta_j}(\bar{x}_j),$$

where  $g_{\theta_i}(\bar{x}_i) = \frac{\sqrt{n_i}}{\sqrt{2\pi}} e^{-\frac{n_i}{2}(\bar{x}_i - \theta_i)^2}$ ,  $\bar{x}_i = \frac{1}{n_i} \sum_{\ell=1}^{n_i} x_{i\ell}$ . Let  $\tau_{ij} = \theta_i - \theta_j$ ,

$1 \leq j \leq k$ ;  $\tau_{ii} = 0$ ,  $1 \leq i \leq k$ , and  $Z_{ij} = \bar{X}_i - \bar{X}_j$ ,  $j \neq i$ . Then for any  $i$ ,

$$\delta_i^{\circ}(z_i) = \begin{cases} 1 & \text{if } z_i \geq d_i, \\ 0 & \text{otherwise,} \end{cases}$$

where  $d_i = (d_{i1}, \dots, d_{i,k-1}, d_{i,k+1}, \dots, d_{ik})$ . Equivalently,

$$(4) \quad \delta_i^{\circ}(\underline{x}) = \begin{cases} 1 & \text{if } \bar{x}_i \geq \max_{j \neq i} (\bar{x}_j + d_{ij}) \\ 0 & \text{otherwise.} \end{cases}$$

We know that

$$P(\bar{X}_i \geq \max_{j \neq i} (\bar{X}_j + d_{ij}))$$

is nondecreasing in  $\theta_i$  and nonincreasing in  $\theta_j$ ,  $j=1, \dots, k$ ,  $j \neq i$ . In this case the, the monotone selection rule also has the above property of monotone behavior in terms of the selection probability. This monotone property is the same as used in the definition of the usual selection procedures. It should be pointed out that when all  $d_{ij}$ 's are negative, the monotone selection procedure  $\delta^{\circ} = (\delta_1^{\circ}, \dots, \delta_k^{\circ})$  given in (4) is the usual Gupta type procedure (cf. [1]) to select a subset containing the best population associated with the largest population associated with the largest  $\theta_i$ 's as follows:

$$\delta_i^{\circ}(\underline{x}) = \begin{cases} 1 & \text{if } \bar{x}_i \geq \max_{1 \leq j \leq k} (\bar{x}_j - (-d_{ij})), \\ 0 & \text{otherwise.} \end{cases}$$

Gupta and Huang [2] have studied the selection rule for the  $k$  normal means problem with a common known variance  $\sigma^2$  based on samples of unequal sizes. In their solution the monotone rules are given by

$$d_{ij} = -d\sigma\sqrt{\frac{1}{n_i} + \frac{1}{n_j}}, \quad d > 0.$$

### References

- [1] Gupta, S. S. (1965). On multiple decision (selection and ranking) rules. Technometrics 7, 225-245.
- [2] Gupta, S. S. and Huang, D. Y. (1976). Subset selection procedures for the means and variances of normal populations: unequal sample sizes case. Sankhyā, 38, Ser. B, 112-128.
- [3] Karlin, S. and Rubin, H. (1956). The theory of decision procedures for distributions with monotone likelihood ratio. Ann. Math. Statist. 27, 272-245.



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Mimeograph Series #78-12 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) An Essentially Complete Class of Multiple Decision Procedures		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) Shanti S. Gupta and Deng-Yuan Huang		6. PERFORMING ORG. REPORT NUMBER Mimeo. Series #78-12
9. PERFORMING ORGANIZATION NAME AND ADDRESS Purdue University Department of Statistics ✓ West Lafayette, IN 47907		8. CONTRACT OR GRANT NUMBER(s) ONR N00014-75-C-0455
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Washington, DC		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June, 1978
		13. NUMBER OF PAGES 7
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Subset selection procedure, monotone likelihood ratio, monotone selection rule, normal means problem, unequal sample sizes. <i>pc-sub<sup>1</sup> pc-sub<sup>k</sup> pc-sub<sup>i</sup> &gt; or = theta sub i</i>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let $\pi_1, \dots, \pi_k$ represent $k$ ( $\geq 2$ ) independent populations. The quality of the $i$ th population $\pi_i$ is characterized by a real-valued parameter $\theta_i$ , usually unknown. We define the best population in terms of a measure of separation between $\theta_i$ 's. A selection of a subset containing the best population is called a correct selection (CS). We restrict attention to rules for which the size of		

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

theta sub i

page

the selected subset is controlled at a given point and the infimum of the probability of correct selection over the parameter space is maximized. The main theorem deals with construction of an essentially complete class of selection rules of the above type. Some classical subset selection rules are shown to belong to this class.

Unclassified